

## Orbital Tori Construction Using Trajectory Following Spectral Methods

By Ralph E. Bordner

Biblioscholar Okt 2012, 2012. Taschenbuch. Book Condition: Neu. 246x189x11 mm. This item is printed on demand - Print on Demand Neuware - The Kolmogorov-Arnold-Moser (KAM) theorem states that lightly perturbed integrable Hamiltonian systems maintain their multiply periodic, toroidal motion in the phase space. The assertion that earth orbiting satellites under the influence of the geopotential mimic this behavior is the underlying premise of this work. This paper focuses on applying trajectory following spectral methods on selected orbits to decompose them into multiperiodic Fourier series, effectively compressing ephemerides for long-term use. The proposed approach focuses on fitting local spectral structures, denoted as frequency clusters, within the sampled orbital data to the analytical form of the windowed, truncated, continuous Fourier transform. This approach is significantly more numerically efficient than fitting every coefficient within the N-tuple Fourier series simultaneously. Numerical results using integrated data show maximum error in orbital torus fits can be kept at the single-digit-meter level per coordinate axis over a 1 year period. 188 pp. Englisch.



## Reviews

It in a single of the most popular ebook. It really is simplified but excitement in the fifty percent from the pdf. It is extremely difficult to leave it before concluding, once you begin to read the book.

-- Joy Langosh

The ideal book i possibly read. It is among the most remarkable pdf i have go through. I am easily could get a enjoyment of reading through a created ebook.

-- Elise Wehner